

The Superior Sticker

An emulsifiable concentrate containing 45% w/w styrene butadiene co-polymer and 10% w/w Alcohol Alkoxylate

FOR USE ONLY AS AN AGRICULTURAL /
HORTICULTURAL / FORESTRY STICKING, EXTENDING
AND WETTING AGENT

ADJ No: 0556 PCS No: 91298

IRRITATING
TO EYES RESPIRATORY SYSTEM
AND SKIN

NET CONTENTS: 5 LITRES @ Protect From Frost



IRRITANT



DE SANGOSSE LTD
HILLSIDE MILL, QUARRY LANE
SWAFFHAM BULBECK • CAMBRIDGE, CB25 0LU
TELEPHONE 01223 811215 • FAX 01223 810020
info@desangosse.co.uk • www.desangosse.co.uk
Distributed in Ireland by T.P. Whelehan Son & Co Ltd.
Finglas Dublin 11, tel 01 8068600

DE SANGOSSE

1 1/10

STATUTORY CONDITIONS RELATING TO USE

FOR USE ONLY AS AN AGRICULTURAL/HORTICULTURAL/FORESTRY STICKING,

EXTENDING AND WETTING AGENT

Maximum Concentration: 0.14% spray solution READ ALL PRECAUTIONS BEFORE USE.

ADJ 0556 PCS 91298

PRECAUTIONS

The (COSHH) Control of Substances Hazardous to Health Regulations may apply to the use of this product at work.

OPERATOR PROTECTION

Engineering control of operator exposure must be used where reasonably practicable in addition to the following personal protective equipment: WEAR SUITABLE PROTECTIVE CLOTHING (COVERALLS),

SUITABLE PROTECTIVE GLOVES AND FACE PROTECTION (FACESHIELD) when handling the concentrate.

However engineering controls may replace personal protective equipment if a COSHH assessment shows that they provide an equal or higher standard of protection.

IRRITATING TO EYES RESPIRATORY SYSTEM AND SKIN

DO NOT BREATHE SPRAY/MISTS.

IN CASE OF CONTACT WITH EYES: rinse immediately with plenty of water and seek medical advice. AVOID CONTACT WITH SKIN

USE IN WELL VENTILATED AREAS

Operators must follow all operator protection instructions for the pesticide. Where the personal protective equipment specified on the adjuvant label is different to that specified on the pesticide label, the highest level of protective equipment must be worn.

ENVIRONMENTAL PROTECTION

DO NOT CONTAMINATE SURFACE WATER OR DITCHES with chemical or used container DO NOT EMPTY INTO DRAINS; dispose of this material and its container in a safe way

STORAGE AND DISPOSAL

KEEP OUT OF REACH OF CHILDREN

KEEP AWAY FROM FOOD. DRINK AND ANIMAL FEEDING STUFFS.

WASH OUT CONTAINER THOROUGHLY, and dispose of safely.

TO AVOID RISKS TO MAN AND THE ENVIRONMENT, COMPLY WITH THE INSTRUCTIONS FOR USE.

RATES OF USE

Bond is used at a rate of 0.1% of the final spray solution in normal conditions and 0.14% in difficult conditions. In practice, due to the UK weather patterns 0.14% is the most commonly used rate.

Water Volume	Required Quantity of BOND	
(litres per hectare)	0.10% Bond	0.14% Bond
70	70 ml	98 ml
100	100 ml	140 ml
200	200 ml	280 ml

MIXING

Half fill the spray tank with water and commence agitation. Add the required quantity of pesticide(s). Top up the spray tank and <u>then</u> add the required quantity of BOND. Continue agitation while topping up the tank and during spraying.

DIRECTIONS FOR USE

BOND is authorised with:-

POTATOES - the <u>full rate</u> of all potato blight fungicides up to their latest approved timing. Edible and Non-edible crops – All approved pesticides which have a recommendation for use with a non ionic wetter.

In addition BOND may be used at a maximum concentration of 0.14% spray solution on the following crops, up to the growth stages specified, with all approved pesticides applied up to their full approved rate.

Crop	Growth Stages before which BOND may be used with approved pesticides
Cereals	up to and including GS 52 (1/4 inflorescence)
Oilseed rape	up to and including 10% potential pods
Broad bean, dwarf French bean, field bean, runner bean	up to and including first pod set
Combining pea and vining pea	up to and including flat pod (non-edible pods only)
Edible podded pea	up to and including pod set
Linseed	up to and including 10% capsules formed
Forage Maize	up to and including tip of tassle visible
Sugar beet	up to and including 6 fully expanded leaves
Potato	up to and including tuber initiation
Brussels Sprout	up to and including lateral buds begin to develop
Cauliflower	up to and including cauliflower heads begin to form; width of growing tip up to 1 cm ²
Apple and pear	up to and including fruit 5-10 mm

In addition to the above conditions BOND is authorised for use with all approved pesticides on all edible crops provided that the statutory conditions of use for the pesticide are followed and that the pesticide is not used at more than 50% of the maximum approved rate for that application.

BOND is also authorised for use with all approved pesticides on all non-edible crops provided that the statutory conditions of use for the pesticide are followed.

BOND may also be used as a spray drift reducing agent where this is a concern. BOND may also be used in tank mixture with trace and macro elements to improve coverage

Do not use in mixes with other adjuvants or if conditions are not recommended for the partner pesticide or trace element.

For Specific Recommendations contact De Sangosse Ltd or its appointed Distributors.

CONDITIONS OF SUPPLY

All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot exercise control over their storage, mixing or use, all conditions and warranties, statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arising from their storage, handling, application or use.