#### **About Corteva Agriscience™**

- A global leader in seed and crop protection created from the former agricultural businesses of Dow AgroSciences, DuPont and Pioneer
- A strong portfolio comprising grassland & maize crop protection, silage inoculants and maize seed
- Corteva's significant investment in innovative science to find and develop new solutions is helping livestock farmers achieve their grassland and forage crop potential

#### \* Ragwort label guidance

Where ragwort is present users should consult the Code of Practice on How to Prevent the Spread of Ragwort. Ragwort plants sprayed with this herbicide are more palatable and contain higher levels of toxins. Animals should be excluded from treated areas until any ragwort has completely recovered or died and there is no visible sign of the dead weed. Do not include treated ragwort in hay or silage crops.

Use plant protection products safely. Always read the label and product information before use. Pay attention to the risk indications and follow the safety precautions on the label.

Triple rinse containers and invert to dry at time of use.

Authorisation Holder: Corteva Agriscience, CPC2 Capital Park, Fulbourn, Cambridge CB21 5XE. Tel: 01462 457272.

Marketing Company: Whelehan Crop Protection, Suite 11/12 Bunkilla Plaza, Bracetown Business Park, Clonee, Co. Meath, Ireland. Tel: 01 4688900 Email: cropprotection@tpwhelehan.ie

®, ™ Trademarks of Corteva Agriscience and its affiliated companies. © 2022 Corteva. Supercedes all previous versions of this leaflet.

Pastor®Trio contains clopyralid, fluroxypyr and florasulam.







**Pastor** Trio is a wider spectrum grassland herbicide giving excellent control of docks, thistles and buttercups. Use on new sown leys, established grassland, grass grown for seed and maize cereals.

See product label for full details.

#### **Corteva Agriscience Forage Apps**

The 'Corteva Forage Farmer App' is packed with new features and resources to help farmers and advisors maximise homegrown forage, including:

- Decision tree tools to help you find the best Corteva and Pioneer product solutions
- Access to newsletters
- Product information
- 'How to' sections and much more.

For more information and for links to download the FREE Corteva Forage Apps, visit corteva.ie/forage or simply scan the QR code.











TPW-2022

**\*** 




For grassland advice call Whelehan Crop Protection on: 01 4688900 or visit: www.corteva.ie/grassland or email: cropprotection@tpwhelehan.ie

grow great grass A



**HERBICIDE** 

# Deadly to weeds, kind to grass.





# Pastor® TRIO

### **HERBICIDE**


Pastor Trio is a wider spectrum grassland herbicide giving excellent control of docks, thistles and buttercups. It is very safe to grass.

#### Control grassland weeds because:

- They compete with grass for space, light, nutrients and water
- It can be used on new sown leys, established grassland, grass grown for seed, maize and cereals
- They are unpalatable to stock
- Thistles can lead to a greater incidence of Orf
- Docks in silage can affect fermentation and reduce quality.

#### **Choose Pastor Trio because it:**

- Delivers excellent control of broadleaved dock, curled dock, creeping thistle, spear thistle, chickweed, creeping and meadow buttercups
- Has a short stock withdrawal period of just 7 days
- Does not hold back grass growth after application.



Where weed populations are below 5%, knapsack control

with Grazon Pro is more cost-effective.

## **Key points:**

•			
Product	Pastor Trio		
Active ingredients	80 g/L clopyralid + 100 g/L fluroxypyr + 2.5 g/L florasulam		
Pack size	3.0 L which treats 3.0 ha (7.4 ac) at 1.0 L/ha and 1.5 ha (3.7 ac) at 2.0 L/ha		
Weeds controlled	Broadleaved dock, curled dock, creeping thistle, spear thistle, chickweed, creeping and meadow buttercups		
Application rate	1.0 L/ha new sown leys, maize, cereals and grass grown for seed 2.0 L/ha established grassland		
Maximum total dose	Same as above		
Water volume	200 L/ha - 400 L/ha for high weed numbers or dense grass swards		
Treatment timing	New sown leys: 1st February to 31st August and from the 3 true leaf stage of the crop Established grassland: 1st February to 30th September Cereals: use from GS13 to before GS39 in Wheat, Barley and before GS32 in Oats  Maize: 10th April to 31st May, between 3 and 6 leaf stage, before the crop is over 20 cm tall and the buttress roots have started to develop on the 1st node		
Weed size Broadleaved and curled dock	Rosette stage, 150-250mm across or high  Docks  Too early  Just right  Too late		
Creeping and spear thistle	Rosette stage, 150-250mm across or high  Thistles  Too early  Too late		
Buttercups	Actively growing, ideally before flowering		
Weed health	Weeds should be actively growing; free from disease or insect damage; not suffering from drought, waterlogging or nutrient deficiency		
Stock exclusion	7 days*		
Cutting interval	Minimum 21 days, ideally 28 days		
Rainfastness	2 hours when applied to a dry leaf		
Clover	Will be damaged; re-introduce after 3 months		

#### **Weeds controlled by Pastor Trio**

Where we have knowledge of how Pastor Trio might affect other grassland weeds we have detailed it in the following tables. These are not recommendations; just an indication of what control might be achieved. Findicates information based on anecdotal or limited data, and as such the user bears the risk in respect of failures concerning efficacy and phytotoxicity.

#### Annual weeds - Pastor Trio at 1.0 L/ha

Bindweed (black)	Fool's parsley	Orache
Bristly ox-tongue	Forget-me-not	Pale persicaria
Charlock (1TL)	Fumitory	Рорру
Chickweed	Groundsel	Redshank
Cleavers	Hemp-nettle	Scarlet pimpernel
Corn chamomile	Himalayan balsam	Shepherd's-purse
Corn marigold	Knotgrass (4TL)	Speedwells
Cranesbill	Mayweeds (1TL)	Spurrey
Dead-nettles	Nettle (small)	Wild radish
Fat-hen (2TL)	Nightshade (black)	Yellow rattle

#### Perennial weeds - Pastor Trio at 2.0 L/ha

Bindweed (field)	Gorse	Old man's beard	
Bracken	Ground elder	Plantain (greater)	
Bramble	Ground ivy	Plantain (ribwort)	
Broom	Hawthorn	Ragwort	
Burdock	Hemlock	Rosebay willowherb	
Buttercups	Hogweed (giant)	Rushes	
Cinquefoil	Horsetail (Equisetum)	Self-heal	
Clover, trefoil	Japanese knotweed	Silverweed	
Coltsfoot	Knapweed (common)	Sorrel (common)	
Cow parsley	Lesser celandine	Thistles	
Daisy (common)	Mallow	Vetch, tare	
Daisy (ox-eye)	Medick	Yarrow	
Dandelion	Mugwort	Yellow/Flag Iris	
Docks	Nettle (common)		
Weed control key	No control		
Good control	No information		
Moderate control	Anecdotal or limited information		

TL = true leaves

Some control